Related edge	Kinematic endpoint
l^+l^- edge	$(m_{ll}^{\max})^2 = (\tilde\xi - \tilde l)(\tilde l - \tilde\chi)/\tilde l$
l^+l^-q edge	$(m_{llq}^{\max})^2 = \begin{cases} \max\left[\frac{(\tilde q-\tilde\xi)(\tilde\xi-\tilde\chi)}{\tilde\xi}, \frac{(\tilde q-\tilde l)(\tilde l-\tilde\chi)}{\tilde l}, \frac{(\tilde q\tilde l-\tilde\xi\tilde\chi)(\tilde\xi-\tilde l)}{\tilde\xi\tilde l}\right] \\ \text{except for the special case in which } \tilde l^2 < \tilde q\tilde\chi < \tilde\xi^2 \\ \text{and } \tilde\xi^2\tilde\chi < \tilde q\tilde l^2 \text{ where one must use } (m_{\tilde q} - m_{\tilde\chi_1^0})^2. \end{cases}$
Xq edge	$(m_{Xq}^{\max})^2 = X + \frac{(\tilde q-\tilde\xi)}{2\tilde\xi}\left[\tilde\xi + X - \tilde\chi + \sqrt{(\tilde\xi - X - \tilde\chi)^2 - 4X\tilde\chi}\right]$
l^+l^-q threshold	$(m_{llq}^{\min})^2 = \begin{cases} [\ 2\tilde l(\tilde q-\tilde\xi)(\tilde\xi-\tilde\chi) + (\tilde q+\tilde\xi)(\tilde\xi-\tilde l)(\tilde l-\tilde\chi) \\ \ -(\tilde q-\tilde\xi)\sqrt{(\tilde\xi+\tilde l)^2(\tilde l+\tilde\chi)^2 - 16\tilde\xi\tilde l^2\tilde\chi}\ \ ]/(4\tilde l\tilde\xi) \end{cases}$
$l_{\text{near}}^{\pm}q$ edge	$(m_{l_{\text{near}}q}^{\max})^2 = (\tilde q - \tilde\xi)(\tilde\xi - \tilde l)/\tilde\xi$ (unobservable)
$l_{\text{far}}^{\pm}q$ edge	$(m_{l_{\text{far}}q}^{\max})^2 = (\tilde q - \tilde\xi)(\tilde l - \tilde\chi)/\tilde l$ (unobservable)
$l^{\pm}q$ high-edge	$(m_{lq(\text{high})}^{\max})^2 = \max\left[(m_{l_{\text{near}}q}^{\max})^2, (m_{l_{\text{far}}q}^{\max})^2\right]$
$l^{\pm}q$ low-edge	$(m_{lq(\text{low})}^{\max})^2 = \min\left[(m_{l_{\text{near}}q}^{\max})^2, (\tilde q - \tilde\xi)(\tilde l - \tilde\chi)/(2\tilde l - \tilde\chi)\right]$
$\tilde l\bar{\tilde l}$ edge	$\Delta M^{\max} = (m_{\tilde l}^2 - m_{\tilde\chi_1^0}^2)/(2m_{\tilde l})$

Table 5.2: *The absolute kinematic endpoints of invariant mass quantities formed from decay chains of the types mentioned in the text for known particle masses. The following shorthand notation has been used:* $\tilde\chi = m_{\tilde\chi_1^0}^2, \tilde l = m_{\tilde l_R}^2, \tilde\xi = m_{\tilde\chi_2^0}^2, \tilde q = m_{\tilde q}^2$ *and* X *is* m_h^2 *or* m_Z^2 *depending on which particle participates in the "branched" decay. The positions of endpoints are derived in Appendix D.*

For completeness, the same table includes the positions of the unobservable near- and far-edges. Note that although the position of the high-edge is just the higher of $m^{\max}_{l_{\rm near}q}$ and $m^{\max}_{l_{\rm far}q}$, the position of the low-edge has a more interesting form. This asymmetry arises because it is not always kinematically possible for the invariant mass of the lepton-quark pair coming from the lower of the near- and far-distribution to approach $\min\left[m^{\max}_{l_{\rm near}q}, m^{\max}_{l_{\rm far}q}\right]$ arbitrarily closely, while simultaneously requiring that this invariant mass is less than that of the other lepton-quark pair.

5.7.6 Dislepton production, $M_{T2}^{\max}(\chi)$, ΔM and the $\tilde{l}\bar{\tilde{l}}$ edge

One of the novel parts of this analysis concerns the introduction of ΔM, a variable designed to better constrain the slepton-LSP mass difference.

It is easiest to describe ΔM in terms of a related variable, $M_{T2}(\chi)$ proposed by the author (in collaboration) in [28].

$M_{T2}(\chi)$ may be used to study events where:

- the event contains two identical two-body decays (i.e. $X \to Y a_1 a_2$ followed by $a_1 \to b_1 c_1$ and $a_2 \to b_2 c_2$),
- $m_A = m_{a_1} \equiv m_{a_2}$ is unknown,
- $m_C = m_{c_1} \equiv m_{c_2}$ is unknown,
- $m_B = m_{b_1} \equiv m_{b_2}$ is known,
- particles of type c are undetectable,
- the longitudinal momentum of the incoming particles is also unknown, and
- Y does not contain any unobservable particles such as neutrinos.

In such cases, the variable provides a kinematic constraint on the masses of particles of type a and type c. In this analysis, $M_{T2}(\chi)$ will be applied to dislepton events of the form $pp \to Y\tilde{l}_R^{\pm}\tilde{l}_R^{\mp} \to Yl^{\pm}l^{\mp}\tilde{\chi}_1^0\tilde{\chi}_1^0$ (Figure 5.4) using the assignment $a = \tilde{l}$, $b = e/\mu$ and $c = \tilde{\chi}_1^0$. These events are used to constrain $m_{\tilde{l}}$ and $m_{\tilde{\chi}_1^0}$ in the following way:

Suppose, for the moment, that the mass of the neutralino were known, and denoted by $m_{\tilde{\chi}}$ (the numbers in $\tilde{\chi}_1^0$ have been dropped to avoid confusion with subscripts indicating which decay the neutralino came from). Then, if it were possible to observe the momenta of each neutralino, and correctly identify its associated lepton, it would be possible to compute the invariant mass of either slepton using the exact relation:

$$m_{\tilde{l}}^2 = m_l^2 + m_{\tilde{\chi}}^2 + 2(E_T^l E_T^{\tilde{\chi}} \cosh(\Delta y) - \mathbf{p}_T^l \cdot \mathbf{p}_T^{\tilde{\chi}}) \tag{5.9}$$

where $E_T^l = \sqrt{{\mathbf{p}_T^l}^2 + m_l^2}$, $E_T^{\tilde{\chi}} = \sqrt{{\mathbf{p}_T^{\tilde{\chi}}}^2 + m_{\tilde{\chi}}^2}$ and Δy is the difference between the rapidities (see Appendix C) of the lepton and the neutralino. This definition includes the lepton masses for completeness, although these are neglected in all computations. Of course, observing the neutralinos is impossible at ATLAS, so the difference in rapidities, Δy, will certainly not be known. Nevertheless, using the fact that $\cosh \Delta y \geq 1$, it is possible to make the reduced claim:

$$m_{\tilde{l}}^2 \geq m_T^2(\mathbf{p}_T^l, \mathbf{p}_T^{\tilde{\chi}}, m_{\tilde{\chi}}) \equiv m_l^2 + m_{\tilde{\chi}}^2 + 2(E_T^l E_T^{\tilde{\chi}} - \mathbf{p}_T^l \cdot \mathbf{p}_T^{\tilde{\chi}}), \tag{5.10}$$

or even the stronger claim:

$$m_{\tilde{l}}^2 \geq \max\{m_T^2(\mathbf{p}_T^{l_1}, \mathbf{p}_T^{\tilde{\chi}_1}, m_{\tilde{\chi}}), m_T^2(\mathbf{p}_T^{l_2}, \mathbf{p}_T^{\tilde{\chi}_2}, m_{\tilde{\chi}})\} \tag{5.11}$$

using both of the decays. But neither $m_T^2(\mathbf{p}_T^l, \mathbf{p}_T^{\tilde{\chi}}, m_{\tilde{\chi}})$ nor the right hand side of (5.11) is an observable quantity, as each relies upon knowledge of the transverse components of the momenta of each neutralino. Under the prevailing assumptions, the only thing known about the neutralinos is the *sum* of their transverse momenta, given by the missing momentum of the event:

$$\not{\mathbf{p}}_T = \mathbf{p}_T^{\tilde{\chi}_1} + \mathbf{p}_T^{\tilde{\chi}_2}. \tag{5.12}$$

Although the form of the splitting (5.12) is not known, it is possible to turn the right hand side of (5.11) into a quantity $M_{T2}^2(m_{\tilde{\chi}})$ which *is* observable, by trying all possible splittings. This motivates the definition:

$$M_{T2}^2(m_{\tilde{\chi}}) \equiv \min_{\not{\mathbf{p}}_1 + \not{\mathbf{p}}_2 = \not{\mathbf{p}}_T} \left[\max \{ m_T^2(\mathbf{p}_T^{l_1}, \not{\mathbf{p}}_1, m_{\tilde{\chi}}), m_T^2(\mathbf{p}_T^{l_2}, \not{\mathbf{p}}_2, m_{\tilde{\chi}}) \} \right], \tag{5.13}$$

which must therefore satisfy

$$m_{\tilde{l}}^2 \geq M_{T2}^2(m_{\tilde{\chi}}). \tag{5.14}$$

In [28], the authors note that it *is* kinematically possible for equality to be reached in (5.14), and that this occurs when $\Delta y_1 = \Delta y_2 = 0$ and $\mathbf{v}_T^{l_1} - \mathbf{v}_T^{\tilde{\chi}_1}$ is parallel (note: not anti-parallel) to $\mathbf{v}_T^{l_2} - \mathbf{v}_T^{\tilde{\chi}_2}$, where $\mathbf{v}_T = \mathbf{p}_T / E_T$. Thus, by construction, $M_{T2}(m_{\tilde{\chi}})$ has the property that, for a large enough number of signal events in a perfect detector,

$$M_{T2}^{\max}(m_{\tilde{\chi}_1^0}) \equiv \max_{\text{events}} \left[M_{T2}(m_{\tilde{\chi}_1^0}) \right] = m_{\tilde{l}}, \tag{5.15}$$

which is equivalent to saying that the endpoint of the $M_{T2}(m_{\tilde{\chi}})$ distribution occurs at the slepton mass.

Note that up to this point it has been necessary to assume knowledge of the neutralino mass, $m_{\tilde{\chi}} \equiv m_{\tilde{\chi}_1^0}$, which is a little unfair, given that it is one of the quantities which the analysis seeks to determine. Formally, then, the M_{T2} distribution is not a single distribution, but a family of distributions: one for each possible value of the neutralino mass. For example; suppose it is proposed that the neutralino mass is equal to "χ" (not necessarily close to the correct value). This would motivate plotting the distribution of $M_{T2}(\chi)$, generated from dislepton events under the hypothesis that the neutralino mass may be taken to be χ rather than $m_{\tilde{\chi}}$ in definition (5.13). Having plotted this distribution, the location $M_{T2}^{\max}(\chi)$ of its endpoint could be ascertained, and the interpretation of $M_{T2}^{\max}(\chi)$ would be "the slepton mass favoured by the data given the assumed neutralino mass".

Thus the overall result obtained from a full set of $M_{T2}(\chi)$ distributions, is a function $M_{T2}^{\max}(\chi)$, describing the most favoured slepton mass as a function of neutralino mass χ.

In Section D.7 of the Appendix it is shown that the function $M_{T2}^{\max}(\chi)$ is in fact dependent on only one unknown parameter $p > 0$; namely the modulus of the three-momentum of the lepton (or the neutralino) in the rest frame of the slepton. In general p is given by

$$p^2 = \frac{m_{\tilde{l}}^4 + m_{\tilde{\chi}_1^0}^4 + m_l^4 - 2(m_{\tilde{l}}^2 m_{\tilde{\chi}_1^0}^2 + m_{\tilde{l}}^2 m_l^2 + m_{\tilde{\chi}_1^0}^2 m_l^2)}{4m_{\tilde{l}}^2}. \tag{5.16}$$

In Section D.7 the general form of the dependence of $M_{T2}^{\max}(\chi)$ upon p is shown to be:

$$M_{T2}^{\max}(\chi) = \sqrt{m_l^2 + p^2} + \sqrt{\chi^2 + p^2}. \tag{5.17}$$

Significantly, (5.17) may be inverted for arbitrary χ yielding:

$$p^2 = \frac{\tilde{L}^4 + \chi^4 + m_l^4 - 2(\tilde{L}^2\chi^2 + \tilde{L}^2 m_l^2 + \chi^2 m_l^2)}{4\tilde{L}^2} = \text{“independent of } \chi\text{”}, \tag{5.18}$$

where $M_{T2}^{\max}(\chi)$ has been abbreviated to $\tilde{L}$. The reader is encouraged to compare (5.18) with (5.16). The result (5.18) is important for two reasons. Firstly it tells us that we no longer need to worry about choosing a particular value of χ in the region of $m_{\tilde{\chi}_1^0}$, and secondly it shows us that p from (5.16) is the quantity which $M_{T2}^{\max}(\chi)$ is *really* constraining. In fact, the most elegant way of extracting p from $M_{T2}(\chi)$ measurements actually *exploits* the freedom to choose an arbitrary value of χ. Paradoxically, the best choice for χ turns out to be neither 0 nor $m_{\tilde{\chi}_1^0}$ (which one might have expected from the original definition of χ) but is in fact m_l. Using this value one may define the quantity $\Delta M \geq 0$ by

$$(\Delta M)^2 \equiv \frac{1}{4}\left(M_{T2}(m_l)\right)^2 - m_l^2 \tag{5.19}$$

and thereby obtain from (5.17) the significant result that:

$$\Delta M^{\max} = p = p(m_{\tilde{l}}, m_{\tilde{\chi}_1^0}, m_l). \tag{5.20}$$

As a consequence of this important result, it is distributions of ΔM which are chosen to be plotted to extract model independent information from dislepton events.[h] The result (5.20) shows us that the location of the ΔM edge (i.e. $\Delta M^{\max}$) is a direct measumenet of p given by (5.16).

Thus far, the only approximation which has been made was the presumption that transverse slepton boosts have a negligible effect on the prediction for $\Delta M^{\max}$ given in (5.20). Within this thesis, however, there is additionally a motivation for neglecting the lepton masses:

In order to maximise the size of the signal, the analysis presented in this thesis will seek both e^+e^- and $\mu^+\mu^-$ events. The presence of the lepton mass in (5.20) (or equivalently in the definition of p) suggests that one ought to analyse events with different

[h]It should be noted that montecarlo studies suggest that the evaluation of $M_{T2}(\chi)$ at values of χ different than $m_{\tilde{\chi}_1^0}$ can introduce a slight dependence of the edge position upon the maximum transverse boost attained by the sleptons. For example, it has been observed that in a situation where *both* sleptons are able to be emitted collinearly and with transverse momenta of 20 GeV (or 100 GeV or 1000 GeV), then the value of $\Delta M^{\max}$ predicted by (5.20) for heavy (left) sleptons at S5 would underestimate th true value by about 2% (or 8% or 14%). As expected, transverse boosts are observed to have no effect on $M_{T2}^{\max}(m_{\tilde{\chi}_1^0})$. The precise reason for this effect is still under investigation, but within this analysis the effect may safely be ignored for two reasons. Firstly, dislepton production is threshold dominated, and so sleptons typically have small transverse momenta. Secondly, and more importantly, the cut limiting jet activity to "at most one with $p_T < 40$ GeV" puts a strong constraint on the momentum available for the sleptons to recoil against. This restricts the washing out of the edge to a 2% effect, which is far less than the order 10% error on the edge position due to statistics. In situations where the effect cannot be neglected, systematic error could be minimised by evaluating $M_{T2}(\chi)$ at the best avaliable guess for $m_{\tilde{\chi}_1^0}$.

leptons differently. Note, however, that in cases where the $\Delta M^{\max}$ is observed to lie at a mass much in excess of m_l, one may immediately deduce that $m_l \ll p$. The assumption $m_l \ll p$, together with definition (5.16), may be used to show that $m_l \ll m_{\tilde{l}} - m_{\tilde{\chi}_1^0}$, which itself permits p to be expanded in terms of $\epsilon = m_l/(m_{\tilde{l}} - m_{\tilde{\chi}_1^0})$ as follows:

$$\Delta M^{\max} = p = p_{\text{approx}} \left\{ 1 - \frac{m_{\tilde{l}}^2 + m_{\tilde{\chi}_1^0}^2}{(m_{\tilde{l}} + m_{\tilde{\chi}_1^0})^2}\epsilon^2 - \frac{2m_{\tilde{l}}^2 m_{\tilde{\chi}_1^0}^2}{(m_{\tilde{l}} + m_{\tilde{\chi}_1^0})^4}\epsilon^4 + O(\epsilon^6) \right\} \tag{5.21}$$

where

$$p_{\text{approx}} \equiv \frac{m_{\tilde{l}}^2 - m_{\tilde{\chi}_1^0}^2}{2m_{\tilde{l}}}. \tag{5.22}$$

This shows that when $\Delta M^{\max} \gg m_l$, then one may take $\Delta M^{\max}$ to be really just a measurement of p_{approx} given by (5.22).

The ΔM distributions arising from two test models examined in this thesis do in fact happen to satisfy $\Delta M^{\max} \gg \max[m_e, m_\mu]$ at the 100 GeV $\gg$ 100 MeV level, and so within this thesis $\Delta M^{\max}$ is taken to be equivalent to p_{approx} as defined in (5.22). This means that within this part of the analysis it is safe to take the leptons to be massless and to analyse both the e^+e^- and the $\mu^+\mu^-$ events together.

To summarise the analysis of the dislepton edge:

- The distributions which will be plotted and whose edges will be identified, will be distributions of ΔM.
- The observation that $\Delta M^{\max} \gg m_\mu$ for the models under study permits us to: neglect lepton masses, analyse both e^+e^- and $\mu^+\mu^-$ events together, and interpret $\Delta M^{\max}$ as a measurement of $p_{\text{approx}} = (m_{\tilde{l}}^2 - m_{\tilde{\chi}_1^0}^2)/(2m_{\tilde{l}})$.
- Were the analysis to be performed at another point where $\Delta M^{\max}$ was much less than (or order) m_l, then the analysis could still proceed, but it would no longer be possible to treat $\Delta M^{\max}$ as a measurement of p_{approx}. $\Delta M^{\max}$ would instead

constrain p. Beacuse p depends on m_l, one would have to consider e^+e^- events separately from $\mu^+\mu^-$ events.

The ΔM edge can thus be included in the analysis to provide an additional constraint on the slepton and neutralino masses. The name 'ΔM' is chosen because, to first order, $\Delta M^{\max} \approx p_{\text{approx}}$ is well approximated by the slepton/neutralino mass difference provided that the slepton is not *much* heavier than the neutralino.

In practice the ΔM edge can be distorted by the finite resolution of the detector or missing energy from soft underlying events. Additionally, standard model (SM) backgrounds provide constraints on minimum detectable slepton/neutralino mass differences (see Section **??**) **XXX FIX REFERENCE!! XXX**. However the most important factor to control is the ability to correctly identify which production processes are contributing to an observed $\Delta M(\chi)$ edge – particularly since it is possible to have multiple edges in the *non* standard model contributions to $\Delta M(\chi)$ distributions. At S5, for example, where both the right- and left-sleptons are lighter than at O1 (see Table 5.1), both $\tilde{l}^+_R\tilde{l}^-_R$ and $\tilde{l}^+_L\tilde{l}^-_L$ events pass the cuts, and two edges are generated. The edge coming from the (lighter) right-slepton falls well under the SM background and so is not measurable, while the (heavier) left-slepton still has a cross section for pair-production high enough to let it form a good edge of its own at $\Delta M(m_{\tilde{\chi}^0_1}) = m_{\tilde{l}_L} - m_{\tilde{\chi}^0_1}$. It is important that this edge is not mistaken for one produced by $\tilde{l}_R$, so methods are needed to dismiss it. Discussion of these issues are delayed until Section **??** XXX FIX REFERENCE!! XXX.

5.7.7 Notes concerning interpretation of endpoint positions

Were all squarks to have the same mass, then (with perfect detector resolution and with infinite statistics) the endpoints of the edges listed above would be found at the positions listed above and summarised in Table 5.2. Note that these positions depend on four unknown parameters, namely the masses of the squark, the slepton and the two neutralinos participating in each decay. The mass of the lightest higgs boson also appears, but it will be assumed that the higgs mass will already be known or will be obtained by other methods (e.g. [14]) to within 2%.

In a more realistic "non-degenerate squark masses" scenario, every distribution with a squark related edge will actually be a superposition of many underlying distributions (one for each squark mass), each having its edge at a slightly different location. This results in some smearing of the edges, even before detector effects (resolutions, acceptances, jet energy calibrations etc.) are taken into account.

It is not possible to measure each of the squark masses separately. Consequently the quantity $m_{\tilde{q}}$ in Table 5.2 will, after being obtained from the "smeared" edges, represent a squark mass *scale* rather than a specific squark mass. In all squark related edges, except the l^+l^-q threshold, the contribution to the outermost part of each edge is provided by the heaviest squarks. In the case of the l^+l^-q threshold, the "true" endpoint is set by the lightest squark. However, if (as at O1 and S5) the other eleven squarks are much heavier (see Table 5.1), it is easier to measure the contribution coming from them. A strong correlation between $m_{\tilde{q}}$ and the mass of the heaviest squark would therefore be expected.

More work would be required to fully understand the "theoretical" systematic errors including the use of a full simulation of the expected edge positions. Better edge models than those used later in this analysis will definitely be needed in order to permit measurements of edge positions to be better correlated with functions of the particle masses. Other sources of systematic errors which require further analysis are the detector effects mentioned above, combinatorial backgrounds near the edges and possible cut biases. Such systematic errors can only meaningfully be studied when real data are available.

It has already been said that, within this thesis, it will be assumed that the above investigations could be performed so as to leave the eventual edge resolutions determined only by statistics and detector resolution. In this analysis, then, all edges are fitted with simple shapes (see Section 5.10.1) with the intention of extracting only an estimate of the *uncertainty* on the edge position, and not to obtain the edge position itself.

5.8 Event generation and pile-up

All events, except those for $q\bar{q} \to W^+W^-$ background processes, were simulated by `HERWIG-6.0` [29]. The W-pair events were generated by `ISAJET-7.42` [30].

At high luminosity, approximately 20 minimum bias events ("pile-up" events) are expected to occur in each bunch crossing. No pile-up events were generated for this analysis, but their effects will have been accounted for within the `ATLFAST` simulation, which alters its reconstruction resolutions in the different luminosity environments appropriately. But resolutions are not the only things which change at high luminosity. For example, the number of jets per bunch crossing with more than a given p_T rises with luminosity, but this effect would not be reproduced by the simulation method described so far. Consequently all cuts which are expected to perform under high luminosity must be checked for susceptibility to the effects which `ATLFAST` does not consider.

Except where stated otherwise, **XXX IS THIS TRUE? XXX,** events were generated and studied in batches corresponding to 100 fb^{-1} in the high luminosity environment. This is expected to correspond to approximately one year of high luminosity running.



Figure 5.7: *Line-shapes used for fits to the mass distributions. Shape (a) (or its reflection in the vertical axis) is the standard "straight line" used to extract the l^+l^-q edge, $l^\pm q$ high-edge, $l^\pm q$ low-edge and l^+l^-q threshold resolutions. Shape (b) models the expected shape of the sharp l^+l^- edge in the absence of detector effects. The l^+l^- edge is actually fitted with shape (c), which is identical to (b) except that it is smeared with a Gaussian resolution whose width is a free parameter of the fit.*

similar quality at O1 (see Figure 5.6: a, b and d). In addition, plots c1 and c2 in Figures 5.5 and 5.6 display the $m_{lq(\text{high})}$ and $m_{lq(\text{low})}$ distributions generated from the modified $l^\pm q$ edge cuts also listed above.

The distributions seen at S5 and O1 are broadly similar to each other. The most obvious difference between the two sets of data is the peak at the Z mass present in Figure 5.6a but absent in Figure 5.5a. This peak comes from direct $\tilde{\chi}_2^0 \to \tilde{\chi}_1^0 Z \to \tilde{\chi}_1^0 l^+ l^-$ decays, which have a branching fraction of 39% at O1 compared with 0.65% at S5. The most common direct neutralino decay mode at S5 is through the h_0 (65%) of which only a tiny proportion (0.02%) decay to the two light leptons as the partial decay width $\Gamma(h_0 \to ff)$ goes approximately as m_f^2. Scenarios in which the l^+l^- edge happens to coincide with the Z peak would require special treatment and are not considered here.

It has already been mentioned that the primary reason for plotting these distributions is not to recover the locations of the edges themelves, but to get an idea of the limit which finite statistics imposes on the resolution of each endpoint. These errors will serve as the foundation for assessing the method as a whole, under the assumption that systematic effects are small in comparison to statistical errors.

In order to extract estimates for the statistical errors on the endpoints, simple fits have been made to the distributions of Figures 5.5 and 5.6. The shapes fitted to the data (see Figure 5.7) and the algorithms for determining the boundaries of the fitted regions have been kept as simple and generic as possible, with the intention of making the fit results both conservative and simple to interpret. Only two shapes were used to fit the edges; a straight line for most edges, and a triangular distribution smeared by a gaussian resolution for the l^+l^- edge. The use of the special method at the l^+l^- edge is justified by the unique step-like nature of its edge (which may not be approximated by a straight line of finite gradient) and also by the fact that the expected shape of the edge is simple to interpret theoretically (see Section 5.7.1) and is unlikely to introduce unwanted model dependence.

To try to avoid bias toward one or either model or toward one or either edge, all the distributions fitted with a *straight line* (ten in total) were fitted by *the same algorithm*. This algorithm was given the tasks of both identifying the range of bins over which to fit, as well as the endpoint of the edge and its gradient. There were a number of inputs to the algorithm common to all distributions, describing "what to look for in an edge" (such as a maximum height which the top part of the edge may reach as a fraction of the height of the bin containing the most events, or a minumum number of events required to be present in each bin over the fit range) but only one input which was permitted to vary from edge to edge; namely the instruction indicating whether to look for a threshold (i.e. left-hand edge) or a forwards edge (i.e. right-hand edge).

The resolutions obtained from the fits to the distributions in Figure 5.5 and Figure 5.6 are listed in Table 5.5.

Given real data, it would be worth putting substantial effort into understanding how detector and physics effects affect the shape of each distribution in turn, particularly in the neighbourhood of each edge. This might significantly improve particle mass resolutions, or at least improve confidence in the errors associated with each edge. But as the events being processed here are entirey synthetic, the fit errors listed in Table 5.5 will be used as they are.

5.10.2 ΔM edge resolution

Determining the likely resolution for the ΔM edge requires a slightly different approach to that used for the "boost invariant" edges above. Whereas the cuts used to obtain the above invariant mass distributions have high SM rejections (primarily due to the presence of at least one high mass $\not{p}_T$ or $M_{\text{effective}}$ requirement), the ΔM cuts cannot be set so hard, primarily because the desired dislepton events have have very little jet activity. (With the dislepton production cross sections typically two orders of magnitude smaller than the squark/gluino production cross sections (Table 5.3) a low efficiency is not affordable.) There are also irreducible SM backgrounds (primarily $W^+W^- \to l^+l^-\nu\bar{\nu}$ and $t\bar{t} \to b\bar{b}W^+W^- \to jjl^+l^-\nu\bar{\nu}$ in cases where jets are below the reconstruction threshold or are outside detector acceptance) which have signatures identical to dislepton events. The smallness of the signal and the presence of these backgrounds would cause problems for naive straight line fitting techniques of the type used above, and so estimation of the edge precision requires the following discussion:

Figure 5.8 shows the ΔM distributions obtained at S5 and O1 after applying the cuts listed in Section 5.9.1 to a 100 fb^{-1} sample of signal dislepton events. Events from lighter sleptons ($\tilde{e}_R$ and $\tilde{\mu}_R$) occupy the unhatched region in each plot, while the events from heavier left-sleptons are cross hatched. It will be noted that events from both light and heavy sleptons succeed in passing the cuts in both models. In principle, then, there are two edges to be observed in each of the models: one for the lighter slepton and one for the heavier. We note, however, that the slepton masses increase, their production is strongly suppressed, and so there are very few heavy slepton events at O1 where there



(a) S5, soft cuts

(b) S5, hard cuts

(c) O1, soft cuts

(d) O1, hard cuts

Figure 5.8: *ΔM distributions obtained at S5 and O1 after applying the cuts listed in Section 5.9.1 to a* $100\ \mathrm{fb}^{-1}$ *sample of $\tilde{e}_R\bar{\tilde{e}}_R$, $\tilde{\mu}_R\bar{\tilde{\mu}}_R$, $\tilde{e}_L\bar{\tilde{e}}_L$ and $\tilde{\mu}_L\bar{\tilde{\mu}}_L$ dislepton events. Events from light right-sleptons (unhatched) are stacked on top of those from heavier left-sleptons (hatched). Only events from OSSF leptons combinations are shown. The plots are generated without OSDF background subtraction, but were it to be performed, no significant differences would be apparent as only 4 (12) signal events are able to pass OSDF soft cuts at S5 (O1). Arrows indicate the values of $\Delta M^{\max}$ predicted by theory for the two types of slepton in each model. A red vertical line is drawn through each plot at half the W mass.*