

Feynman's Golden Rule

$$\Gamma_{ji} = 2\pi |\mathcal{T}_{ji}|^2 \rho(E_j)$$

$$\mathcal{T}_{ji} = \langle j | \hat{H}_{int} | i \rangle + \sum_{l \neq i} \frac{\langle j | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | i \rangle}{E_i - E_j} + \dots$$

Flux + Phase-space connection to cross sections σ , $\frac{d\sigma}{dt}$ etc.

⊕

"Old Fashioned Perturbation Theory"

Issue

100% Valid ✓ But calculating $\mathcal{T}_{ji}$ is not convenient.

One source of big problems is normalisation:

$$\langle \psi | \psi \rangle = 1 \text{ (or const.)}$$

(Volumes length contract by factor γ under BOOST)

↓

Part of resolution is state renormalisation:

Single particle state:

$$|\psi'\rangle = \sqrt{2E} |\psi\rangle$$

so that $\int \langle \psi' | \psi' \rangle dV = 2E \int \langle \psi | \psi \rangle dV = 2E = \gamma \times \text{const}$

Multi-particle state:

$$|a', b', c'\rangle = \sqrt{2E_a} \sqrt{2E_b} \sqrt{2E_c} |a, b, c\rangle$$

Turns out to make $\langle \psi'_1 \psi'_2 \dots | \hat{H}_{int} | \dots \psi'_n \rangle$ Lorentz invariant see why later.

↓


Motivates this definition

$$\frac{M_{abc \dots de}}{\sqrt{2E_a} \sqrt{2E_b} \dots \sqrt{2E_e}} = \frac{\langle abc \dots | \hat{H}_{int} | \dots de \rangle}{\text{CONSEQUENCE}}$$

...only later see why this is useful.

Consider $\mathcal{T}_{ji} = \langle j | \hat{H}_{int} | i \rangle + \sum_{l \neq i} \frac{\langle j | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | i \rangle}{E_i - E_j} + \dots$

(a = shorthand for p_a)

For special case of  Mom Cons $\Rightarrow a + b = c + d$

" $|i\rangle = |a, b\rangle$ "
" $|j\rangle = |c, d\rangle$ "

(Shot states $|i\rangle$ are being summed over?)


Every basis state of $\hat{H}_{free}$:

$$|j\rangle \in \{ |0\rangle, |a, b\rangle, |c, d\rangle, |c, d\rangle, |a, b\rangle, \dots, |a, b, c, d\rangle, |E_{kin} \text{ Messy}\rangle, |j_a\rangle, |j_b\rangle, \dots \}$$

In this giant list, consider just two special entries:

" $|j_a\rangle = |c, b, \underline{z} = \underline{a} - \underline{c}\rangle$ $\underline{z}$ = "particle with mass m & mom $\underline{z}$ "
" $|j_b\rangle = |a, d, \underline{y} = \underline{b} - \underline{d}\rangle$ $\underline{y}$ = "particle with mass m & mom $\underline{y}$ "

Notes:

- (mom cons $p_{initial} = p_{final} \Rightarrow a + b = c + d \Rightarrow \underline{z} = -\underline{y}$)
 $\therefore \underline{z} \text{ \& } \underline{y}$ are heading in "opposite directions",
 and $E_x = \sqrt{m^2 + |\underline{z}|^2} = \sqrt{m^2 + |\underline{y}|^2} = E_y$,
- $\underline{x} \text{ \& } \underline{y}$ have correct 3-momenta to conserve momentum in the following ways:

- 4-momentum is not conserved, though!
 (i.e. $\nexists E_i, E_j, E_c + E_d = E_a + E_b$
 and $E_i, E_j, E_a + E_b = E_c + E_d$)
 then $(a^0 + b^0 = c^0 + d^0) \Rightarrow (E_a + E_b = E_c + E_d)$
 but $(a^0 + b^0 = c^0 + d^0) \not\Rightarrow (E_a + E_b = E_i)$
 and $(a^0 + b^0 = c^0 + d^0) \not\Rightarrow (E_a + E_b = E_j)$.

A feel for terms in the interaction Hamiltonian

$$\hat{H}_{int} = e \hat{\phi} \hat{\psi} \hat{\psi} + e \hat{\phi} \hat{\psi} \hat{\psi} + \dots$$

$$\hat{\phi} \sim \int \hat{a}_p + \hat{a}_p^\dagger dp$$

$$\hat{\psi} \sim \int \hat{b}_p + \hat{b}_p^\dagger dp$$

$\therefore$

$\hat{\phi} |0\rangle \sim |0\rangle$
 $\hat{\phi} \hat{\phi} |0\rangle \sim |0\rangle + \int |p, q\rangle dp dq$
 $\hat{\phi} \hat{\phi} \hat{\phi} |0\rangle \sim |0\rangle + \int |p\rangle dp + \int |p, q, r\rangle dp dq dr$
 $\hat{\phi} \hat{\phi} \hat{\phi} \hat{\phi} |0\rangle \sim |0\rangle + \int |p, q\rangle dp dq + \int |p, q, r, s\rangle dp dq dr ds$

Since " $|i\rangle = |a, b\rangle$ " and " $|j\rangle = |c, d\rangle$ " consider the following M_{ji} :

$$M_{ji} = \sqrt{2E_a} \sqrt{2E_b} \sqrt{2E_c} \sqrt{2E_d} \mathcal{T}_{ji}$$

$$= \sqrt{2E_a} \sqrt{2E_b} \sqrt{2E_c} \sqrt{2E_d} \left(\langle c, d | \hat{H}_{int} | a, b \rangle + \sum_{l \neq i} \frac{\langle c, d | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | a, b \rangle}{(E_a + E_b) - E_j} + \dots \right)$$

$$= \langle c', d' | \hat{H}_{int} | a', b' \rangle + \sum_{l \neq i} \frac{\langle c', d' | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - E_j} + \dots$$

$$= M_{cd;ab} + \sum_{l \neq i} \frac{\langle c', d' | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - E_j} + \dots$$

$$= M_{cd;ab} + \frac{\langle c', d' | \hat{H}_{int} | j_a \rangle \langle j_a | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - E_{j_a}} \leftarrow j_a \text{ term in sum}$$

$$+ \frac{\langle c', d' | \hat{H}_{int} | j_b \rangle \langle j_b | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - E_{j_b}} \leftarrow j_b \text{ term in sum}$$

$$+ \sum_{l \neq \{j_a, j_b, |0\rangle, |j_a\rangle, |j_b\rangle\}} \frac{\langle c', d' | \hat{H}_{int} | l \rangle \langle l | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - E_j} \leftarrow \text{all other terms in sum}$$

+ ... $\leftarrow$ all other terms in other sums.

call all of this "..." in next line

$$= M_{cd;ab} + \frac{\langle c', d' | \hat{H}_{int} | c, b, \underline{z} \rangle \langle c, b, \underline{z} | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - (E_c + E_b + E_z)} + \dots$$

[If momenta were not discretised (not finite box) we would need same kind of $\int d^3c d^3b d^3z$ here, but we gloss over such details.]

$$+ \frac{\langle c', d' | \hat{H}_{int} | a, d, \underline{y} \rangle \langle a, d, \underline{y} | \hat{H}_{int} | a', b' \rangle}{(E_a + E_b) - (E_a + E_d + E_y)} + \dots$$

$$= M_{cd;ab} + \frac{\langle c', d' | \hat{H}_{int} | c, b, \underline{z} \rangle \langle c, b, \underline{z} | \hat{H}_{int} | a', b' \rangle}{E_a - E_c - E_x} \cdot \frac{1}{2E_x \sqrt{2E_b} \sqrt{2E_x}}$$

$$+ \frac{\langle c', d' | \hat{H}_{int} | a, d, \underline{y} \rangle \langle a, d, \underline{y} | \hat{H}_{int} | a', b' \rangle}{E_b - E_d - E_y} \cdot \frac{1}{2E_x \sqrt{2E_x} \sqrt{2E_y}} + \dots$$

[Cancellations shown rely on $\hat{H}_{int}$ having certain undiscussed properties. E.g. $\hat{H}_{int} = \hat{\phi} \hat{\psi} \hat{\psi}$ (a) ψ -type field (b) ϕ -type field]

$$= M_{cd;ab} + \frac{\langle d' | \hat{H}_{int} | b, \underline{z} \rangle \langle c, \underline{z} | \hat{H}_{int} | a' \rangle}{E_a - E_c - E_x} \cdot \frac{1}{2E_x}$$

$$+ \frac{\langle c' | \hat{H}_{int} | a, \underline{y} \rangle \langle d, \underline{y} | \hat{H}_{int} | b' \rangle}{E_b - E_d - E_y} \cdot \frac{1}{2E_y} + \dots$$

$$= M_{cd;ab} + \frac{M_{d;bx} M_{cx;a}}{E_a - E_c - E_x} \cdot \frac{1}{2E_x}$$

$$+ \frac{M_{c;ay} M_{dy;b}}{E_c - E_a - E_y} \cdot \frac{1}{2E_y} + \dots$$

[Since $(E_a + E_b = E_c + E_d) \Rightarrow (E_b - E_d = E_c - E_a)$]

$$\sim h \delta_{\underline{c}+\underline{b}, \underline{a}+\underline{d}} + \frac{g_{cd;bx} g_{cx;a}}{(E_a - E_c - E_x) 2E_x} \leftarrow \text{[If momenta are discrete (ie finite box) then } \delta_{\underline{a}+\underline{b}, \underline{c}+\underline{d}} \leftarrow \delta_{\underline{a}+\underline{b}, \underline{c}+\underline{d}} \text{ no four-point interaction} \right.$$

$$+ \frac{g_{cd;ay} g_{dy;b}}{(E_c - E_a - E_y) 2E_y} + \dots$$

Continuous case would contain Dirac delta-functions ...
 $\int M_{d;bx} d^3x \sim g \int \delta^3(\underline{d} - (\underline{b} + \underline{x})) d^3x$ etc
 (Such δ -funes would actually also ensure $\underline{z} = -\underline{y}$, so no need to separate $|j_a\rangle$ and $|j_b\rangle$ either!)

$$= h \cdot 1 + \frac{g_{cd;bx} \cdot 1}{(E_a - E_c - E_x) 2E_x}$$

$$- \frac{g_{cd;ay} \cdot 1}{(-E_c + E_a + E_x) 2E_x} + \dots$$

However - as this is only sketch such details are ignored. See QFT & AQFT courses for full details!

$$= h + \frac{g_{cd;bx} [(E_a - E_c) + E_x] - [(E_a - E_c) - E_x]}{2E_x ((E_a - E_c) - E_x)((E_a - E_c) + E_x)} + \dots$$

$$= h + \frac{g_{cd;bx}}{(E_a - E_c)^2 - E_x^2} + \dots$$


So now, and only now Define a quantity q^μ by:

$$\boxed{q^\mu \equiv a^\mu - c^\mu} \quad (4\text{-mom conserving}) \quad (a^\mu + b^\mu = c^\mu + d^\mu \Rightarrow q^\mu = d^\mu - b^\mu \text{ too})$$

Clearly $q^\mu = \begin{pmatrix} E_a - E_c \\ \underline{a} - \underline{c} \end{pmatrix} \therefore q^2 = q^\mu q_\mu = (E_a - E_c)^2 - |\underline{a} - \underline{c}|^2$

$$\therefore M_{ji} \sim h + \frac{g_{cd;bx}}{q^2 + |\underline{a} - \underline{c}|^2 - E_x^2} + \dots$$

$$= h + \frac{g_{cd;bx}}{q^2 + |\underline{a} - \underline{c}|^2 - (m^2 + |\underline{z}|^2)} + \dots$$

$$= h + \frac{g_{cd;bx}}{q^2 - m^2} \quad \square$$


could add structure of $\hat{H}_{int}$ to explain 3 & 4-point interactions