


Dr C.G. Lester, 2023




H10: Leptonic Weak Interactions and Neutrino Deep Inelastic Scattering

Aside : Neutrino Flavours

- Recent experiments (see Handout 11) imply neutrinos have mass (albeit very small).
- The textbook neutrino states, ν_e, ν_μ, ν_τ , are not the fundamental particles; these are ν_1, ν_2, ν_3 .
- Concepts like ‘electron number’ conservation are now known **not** to hold.
- So what are ν_e, ν_μ, ν_τ ?
- Never directly observe neutrinos — can only detect them by their weak interactions. Hence **by definition** ν_e is the neutrino state produced along with an electron. Similarly, charged current weak interactions of the state ν_e produce an electron.




$$\{\nu_e, \nu_\mu, \nu_\tau\} = \text{weak eigenstates}$$

- Unless dealing with very large distances: the neutrino produced with a positron will interact to produce an electron. For the discussion of the weak interaction continue to use ν_e, ν_μ, ν_τ as if they were the fundamental particle states.




Muon Decay and Lepton Universality

- The leptonic charged current (W^\pm) interaction vertices are:



- Consider muon decay:



- It is straight-forward to write down the matrix element




$$M_{fi} = \frac{g_W^{(e)} g_W^{(\mu)}}{8m_W^2} [\bar{u}(p_3)\gamma^\mu(1 - \gamma^5)u(p_1)] g_{\mu\nu} [\bar{u}(p_2)\gamma^\nu(1 - \gamma^5)v(p_4)].$$

- Note: for lepton decay $q^2 \ll m_W^2$ so propagator is a constant (Fermi theory limit!).
- Its evaluation and subsequent treatment of a three-body decay is rather tricky (and not particularly interesting). Here will simply quote the result (over page):

- The muon to electron rate is

$$\Gamma(\mu \rightarrow e\nu\nu) = \frac{G_F^e G_F^\mu m_\mu^5}{192\pi^3} = \frac{1}{\tau_\mu} \quad \text{with} \quad G_F = \frac{g_W^2}{4\sqrt{2}m_W^2}.$$

- Similarly for tau to electron $\Gamma(\tau \rightarrow e\nu\nu) = \frac{G_F^e G_F^\tau m_\tau^5}{192\pi^3}$ however, the tau can decay to a number of final states:



$$Br(\tau \rightarrow e\nu\nu) = 0.1784(5) \quad Br(\tau \rightarrow \mu\nu\nu) = 0.1736(5)$$

- Recall total width (total transition rate) is the sum of the partial widths

$$\Gamma = \sum_i \Gamma_i = \frac{1}{\tau}$$

- Can relate partial decay width to total decay width and therefore lifetime:

$$\Gamma(\tau \rightarrow e\nu\nu) = \Gamma_\tau Br(\tau \rightarrow e\nu\nu) = Br(\tau \rightarrow e\nu\nu)/\tau_\tau$$

- Therefore predict $\tau_\mu = \frac{192\pi^3}{G_F^e G_F^\mu m_\mu^5}$ $\tau_\tau = \frac{192\pi^3}{G_F^e G_F^\tau m_\tau^5} Br(\tau \rightarrow e\nu\nu)$.

- All these quantities are precisely measured:


$$\left. \begin{array}{l} m_\mu = 0.1056583692(94) \text{ GeV} \\ m_\tau = 1.77699(28) \text{ GeV} \\ \tau_\tau = 0.2906(10) \times 10^{-12} \text{ s} \\ \tau_\mu = 2.19703(4) \times 10^{-6} \text{ s} \\ Br(\tau \rightarrow e\nu\nu) = 0.1784(5) \end{array} \right\} \text{ so}$$

$$\frac{G_F^\tau}{G_F^\mu} = \frac{m_\mu^5 \tau_\mu}{m_\tau^5 \tau_\tau} Br(\tau \rightarrow e\nu\nu) = 1.0024 \pm 0.0033.$$

- Similarly by comparing $Br(\tau \rightarrow \mu\nu\nu)$ and $Br(\tau \rightarrow e\nu\nu)$

$$\frac{G_F^e}{G_F^\mu} = 1.000 \pm 0.004.$$

The above demonstrates the weak charged current is the same for all leptonic vertices. This is referred to as 'Charged Current Lepton Universality'.




Neutrino Scattering

In Handout 6 considered electron-proton Deep Inelastic Scattering where a virtual photon is used to probe nucleon structure.

- Can also consider the weak interaction equivalent: Neutrino Deep Inelastic Scattering where a virtual W -boson probes the structure of nucleons. **This provides additional information about parton structure functions.**
- provides a good example of calculations of weak interaction cross sections.



Neutrino beams are needed. To make them:

- Smash high energy protons into a fixed target to make hadrons.
- Focus positive pions/kaons.
- Allow them to decay $\pi^+ \rightarrow \mu^+ \nu_\mu + K^+ \rightarrow \mu^+ \nu_\mu$ ($BR \approx 64\%$);
- this gives a beam of “collimated” ν_μ .
- Focus negative pions/kaons to give beam of $\bar{\nu}_\mu$.



Neutrino-Quark Scattering

- For ν_μ -proton Deep Inelastic Scattering the underlying process is $\nu_\mu d \rightarrow \mu^- u$



- In the limit $q^2 \ll m_W^2$ the W -boson propagator is $\approx ig_{\mu\nu}/m_W^2$. so the Feynman rules give:

$$-iM_{fi} = \left[-i \frac{g_W}{\sqrt{2}} \bar{u}(p_3) \gamma^\mu \frac{1}{2} (1 - \gamma^5) u(p_1) \right] \frac{ig_{\mu\nu}}{m_W^2} \left[-i \frac{g_W}{\sqrt{2}} \bar{u}(p_4) \frac{1}{2} \gamma^\nu (1 - \gamma^5) u(p_2) \right]$$

so

$$M_{fi} = \frac{g_W^2}{2m_W^2} g_{\mu\nu} \left[\bar{u}(p_3) \gamma^\mu \frac{1}{2} (1 - \gamma^5) u(p_1) \right] \left[\bar{u}(p_4) \frac{1}{2} \gamma^\nu (1 - \gamma^5) u(p_2) \right].$$

- Evaluate the matrix element in the extreme relativistic limit where the muon and quark masses can be neglected.

- In this limit the helicity states are equivalent to the chiral states. Furthermore

$$\frac{1}{2}(1 - \gamma^5)u_{\uparrow}(p_1) = 0 \quad \text{and} \quad \frac{1}{2}(1 - \gamma^5)u_{\downarrow}(p_1) = u_{\downarrow}(p_1)$$


so $M_{fi} = 0$ for $u_{\uparrow}(p_1)$ and $u_{\uparrow}(p_2)$.

- Since the weak interaction 'conserves the helicity', the only helicity combination where the matrix element is non-zero is

$$M_{fi} = \frac{g_W^2}{2m_W^2} g_{\mu\nu} [\bar{u}_{\downarrow}(p_3)\gamma^{\mu}u_{\downarrow}(p_1)] [\bar{u}_{\downarrow}(p_4)\gamma^{\nu}u_{\downarrow}(p_2)].$$


(We could have written this down straight away as in the ultra-relativistic limit only LH helicity particle states participate in the weak interaction.)

- We will next consider this scattering in the C.o.M frame:



Evaluation of Neutrino-Quark Scattering Matrix Element

- Go through the calculation in gory detail (fortunately only one helicity combination)
- In the $\nu_\mu d$ CMS frame, neglecting particle masses:



$$p_1 = (E, 0, 0, E),$$

$$p_2 = (E, 0, 0, -E)$$

$$p_3 = (E, E \sin \theta^*, 0, E \cos \theta^*)$$

$$p_4 = (E, -E \sin \theta^*, 0, -E \cos \theta^*)$$

- Dealing with LH helicity particle spinors. From Handout 2 (page 106), for a massless particle travelling in direction θ, ϕ :

$$u_\downarrow = \sqrt{E} \begin{pmatrix} -s \\ ce^{i\phi} \\ s \\ -ce^{i\phi} \end{pmatrix} \quad c = \cos \frac{\theta}{2}; \quad s = \sin \frac{\theta}{2}$$

- Here $(\theta_1, \phi_1) = (0, 0)$; $(\theta_2, \phi_2) = (\pi, 0)$; $(\theta_3, \phi_3) = (\theta^*, 0)$; $(\theta_4, \phi_4) = (\pi - \theta^*, \pi)$ giving:

$$u_\downarrow(p_1) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}; \quad u_\downarrow(p_2) = \sqrt{E} \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}; \quad u_\downarrow(p_3) = \sqrt{E} \begin{pmatrix} -s \\ c \\ s \\ -c \end{pmatrix}; \quad u_\downarrow(p_4) = \sqrt{E} \begin{pmatrix} -c \\ -s \\ c \\ s \end{pmatrix}.$$

Neutrino-Quark scattering (cont.)

- To calculate

$$M_{fi} = \frac{g_W^2}{2m_W^2} g_{\mu\nu} [\bar{u}_\downarrow(p_3)\gamma^\mu u_\downarrow(p_1)] [\bar{u}_\downarrow(p_4)\gamma^\nu u_\downarrow(p_2)]$$

need twice to evaluate terms of the form

$$\begin{aligned} \bar{\psi}\gamma^0\phi &= \psi^\dagger\gamma^0\gamma^0\phi = \psi_1^*\phi_1 + \psi_2^*\phi_2 + \psi_3^*\phi_3 + \psi_4^*\phi_4, \\ \bar{\psi}\gamma^1\phi &= \psi^\dagger\gamma^0\gamma^1\phi = \psi_1^*\phi_4 + \psi_2^*\phi_3 + \psi_3^*\phi_2 + \psi_4^*\phi_1, \\ \bar{\psi}\gamma^2\phi &= \psi^\dagger\gamma^0\gamma^2\phi = -i(\psi_1^*\phi_4 - \psi_2^*\phi_3 + \psi_3^*\phi_2 - \psi_4^*\phi_1), \\ \bar{\psi}\gamma^3\phi &= \psi^\dagger\gamma^0\gamma^3\phi = \psi_1^*\phi_3 - \psi_2^*\phi_4 + \psi_3^*\phi_1 - \psi_4^*\phi_2. \end{aligned}$$

- Using

$$u_\downarrow(p_1) = \sqrt{E} \begin{pmatrix} 0 \\ 1 \\ 0 \\ -1 \end{pmatrix}; \quad u_\downarrow(p_2) = \sqrt{E} \begin{pmatrix} -1 \\ 0 \\ 1 \\ 0 \end{pmatrix}; \quad u_\downarrow(p_3) = \sqrt{E} \begin{pmatrix} -s \\ c \\ s \\ -c \end{pmatrix}; \quad u_\downarrow(p_4) = \sqrt{E} \begin{pmatrix} -c \\ -s \\ c \\ s \end{pmatrix}$$

we get

$$\begin{aligned} \bar{u}_\downarrow(p_3)\gamma^\mu u_\downarrow(p_1) &= 2E(c, s, -is, c) \\ \bar{u}_\downarrow(p_4)\gamma^\nu u_\downarrow(p_2) &= 2E(c, -s, -is, -c) \end{aligned}$$

$$\Rightarrow M_{fi} = \frac{g_W^2}{2m_W^2} 4E^2(c^2 + s^2 + s^2 + c^2) = \frac{g_W^2 \hat{s}}{m_W^2}$$


where $\hat{s} = (2E)^2$.

Neutrino-Quark scattering (cont.)

Note that the Matrix Element is isotropic:

$$M_{fi} = \frac{g_W^2}{m_W^2} \hat{s}.$$

We could have anticipated this since the helicity combination (spins anti-parallel) has $S_z = 0$ so expect no preferred polar angle.



- As before need to sum over all possible spin states and average over all possible initial state spin states. Here only one possible spin combination ($LL \rightarrow LL$) and only two possible initial state combinations (the neutrino is always produced in a LH helicity state).

$$\langle |M_{fi}|^2 \rangle = \frac{1}{2} \left| \frac{g_W^2}{m_W^2} \hat{s} \right|^2$$

The factor of a half arises because half of the time the quark will be in a RH states and won't participate in the charged current Weak interaction.

- From Handout 1, in the extreme relativistic limit, the cross section for any $2 \rightarrow 2$ body scattering process is

$$\frac{d\sigma}{d\Omega^*} = \frac{1}{64\pi^2 \hat{s}} \langle |M_{fi}|^2 \rangle.$$

Neutrino-Quark scattering (conclusion).

- Therefore:

$$\frac{d\sigma}{d\Omega^*} = \frac{1}{64\pi^2\hat{s}} \langle |M_{fi}|^2 \rangle = \frac{1}{64\pi\hat{s}} \frac{1}{2} \left(\frac{g_W^2 \hat{s}}{m_W^2} \right)^2 = \left(\frac{g_W^2}{8\sqrt{2}\pi m_W^2} \right)^2 \hat{s}.$$

- Using $\frac{G_F}{\sqrt{2}} = \frac{g_W^2}{8m_W^2}$ the above simplifies to


$$\boxed{\frac{d\sigma}{d\Omega^*} = \frac{G_F^2}{4\pi^2} \hat{s}} \quad (127)$$

and integrating this isotropic distribution over $d\Omega^*$ gives

$$\boxed{\sigma_{\nu q} = \frac{G_F^2 \hat{s}}{\pi}} \quad (128)$$

- Since the cross section is a (longitudinally) Lorentz invariant, (128) is also the cross section for scattering in the lab frame.

Antineutrino-Quark Scattering





- In the ultra-relativistic limit, the charged-current interaction matrix element is:

$$M_{fi} = \frac{g_W^2}{2m_W^2} g_{\mu\nu} \left[\bar{v}(p_1) \gamma^\mu \frac{1}{2}(1 - \gamma^5) v(p_3) \right] \left[\bar{u}(p_4) \gamma^\nu \frac{1}{2}(1 - \gamma^5) u(p_2) \right]$$

- In the extreme relativistic limit only **LH Helicity particles** and **RH Helicity anti-particles** participate in the charged current weak interaction:

$$M_{fi} = \frac{g_W^2}{2m_W^2} g_{\mu\nu} [\bar{v}_\uparrow(p_1) \gamma^\mu v_\uparrow(p_3)] [\bar{u}_\downarrow(p_4) \gamma^\nu u_\downarrow(p_2)]$$

In terms of the particle spins it can be seen that the interaction occurs in a total angular momentum $J = 1$ state:



Antineutrino-Quark Scattering (cont.)

- Similarly to the neutrino-quark scattering calculation obtain:

$$\frac{d\sigma_{\bar{\nu}q}}{d\Omega^*} = \frac{d\sigma_{\nu q}}{d\Omega^*} \frac{1}{4} (1 + \cos\theta^*)^2 \hat{s}$$


- The factor $\frac{1}{4}(1 + \cos\theta^*)^2$ can be understood in terms of the overlap of the initial and final angular momentum wave-functions.
- Integrating over solid angle:

$$\int (1 + \cos\theta^*)^2 d\Omega^* = 2\pi \int_{-1}^{+1} (1 + \cos\theta^*)^2 d(\cos\theta^*) = \frac{16\pi}{3}$$

$$\Rightarrow \boxed{\sigma_{\bar{\nu}q} = \frac{G_F^2 \hat{s}}{3\pi}}$$

- This is a factor three smaller than the neutrino quark cross-section:

$$\boxed{\frac{\sigma_{\bar{\nu}q}}{\sigma_{\nu q}} = \frac{1}{3}}$$




(Anti)neutrino-(Anti)quark Scattering

- Non-zero anti-quark component to the nucleon also consider scattering from q
- Cross-sections can be obtained immediately by comparing with quark scattering and remembering to only include **LH particles** and **RH anti-particles**

$S_z = 0$	$S_z = +1$	$S_z = -1$	$S_z = 0$
$\frac{d\sigma_{\nu q}}{d\Omega^*} = \frac{G_F^2}{4\pi^2} \hat{s}$	$\frac{d\sigma_{\bar{\nu} q}}{d\Omega^*} = \frac{G_F^2}{16\pi^2} (1 + \cos\theta^*)^2 \hat{s}$	$\frac{d\sigma_{\nu \bar{q}}}{d\Omega^*} = \frac{G_F^2}{16\pi^2} (1 + \cos\theta^*)^2 \hat{s}$	$\frac{d\sigma_{\bar{\nu} \bar{q}}}{d\Omega^*} = \frac{G_F^2}{4\pi^2} \hat{s}$
$\sigma_{\nu q} = \frac{G_F^2 \hat{s}}{\pi}$	$\sigma_{\bar{\nu} q} = \frac{G_F^2 \hat{s}}{3\pi}$	$\sigma_{\nu \bar{q}} = \frac{G_F^2 \hat{s}}{3\pi}$	$\sigma_{\bar{\nu} \bar{q}} = \frac{G_F^2 \hat{s}}{\pi}$

Differential Cross Section in form $d\sigma/dy$

- We derived the differential neutrino scattering cross sections in C.o.M frame. Let us convert them to Lorentz invariant form.



- As for DIS use Lorentz invariant $y \equiv \frac{p_2 \cdot q}{p_2 \cdot p_1}$.
- In relativistic limit y can be expressed in terms of the C.o.M. scattering angle: $y = \frac{1}{2}(1 - \cos \theta^*)$.
- In the lab. frame $y = 1 - \frac{E_3}{E_1}$.

- Convert from $\frac{d\sigma}{d\Omega^*} \rightarrow \frac{d\sigma}{dy}$ using

$$\frac{d\sigma}{dy} = \left| \frac{d\cos\theta^*}{dy} \right| \frac{d\sigma}{d\cos\theta^*} = \left| \frac{d\cos\theta^*}{dy} \right| 2\pi \frac{d\sigma}{d\Omega^*} = 4\pi \frac{d\sigma}{d\Omega^*}.$$

- But we already found (in (127)) that: $\frac{d\sigma}{d\Omega^*} = \frac{G_F^2}{4\pi^2} \hat{s}$ hence:

$$\boxed{\frac{d\sigma_{\nu q}}{dy} = \frac{d\sigma_{\bar{\nu} \bar{q}}}{dy} = \frac{G_F^2}{\pi} \hat{s}}.$$

Similarly

$$\frac{d\sigma_{\bar{\nu}q}}{d\Omega^*} = \frac{d\sigma_{\nu\bar{q}}}{d\Omega^*} = \frac{G_F^2}{16\pi^2} (1 + \cos\theta^*)^2 \hat{s}$$

becomes

$$\frac{d\sigma_{\bar{\nu}q}}{d\Omega^*} = \frac{d\sigma_{\nu\bar{q}}}{d\Omega^*} = \frac{G_F^2}{16\pi^2} (1 + \cos\theta^*)^2 \hat{s}$$

from $y = \frac{1}{2}(1 - \cos\theta^*) \rightarrow 1 - y = \frac{1}{2}(1 + \cos\theta^*)$ and hence

$$\boxed{\frac{d\sigma_{\bar{\nu}q}}{dy} = \frac{d\sigma_{\nu\bar{q}}}{dy} = \frac{G_F^2}{\pi} (1 - y)^2 \hat{s}}$$


For comparison, the electro-magnetic $e^\pm q \rightarrow e^\pm q$ cross section is:

$$\text{QED:} \quad \frac{d\sigma_{e^\pm q}}{dy} = \frac{2\pi\alpha^2}{q^4} e_q^2 \left[1 + (1 - y)^2 \right] \hat{s}$$


$$\text{Weak:} \quad \frac{d\sigma_{\bar{\nu}q}}{dy} = \frac{d\sigma_{\nu\bar{q}}}{dy} = \frac{G_F^2}{\pi} (1 - y)^2 \hat{s}$$

(Differences are in (i) **Helicity Structure**, and (ii) in **interaction+propagator**.)

Parton Model For Neutrino Deep Inelastic Scattering



Scattering from a proton with structure functions



Scattering from a point-like quark within the proton

- Neutrino-proton scattering can occur via scattering from a **down-quark** or from an **up-antiquark**.
- In the parton model, number of down quarks within the proton in the $x \rightarrow x + dx$ momentum fraction range is $d^P(x) dx$. Their contribution to the neutrino scattering cross-section is obtained by multiplying by the $\nu_\mu d \rightarrow \mu^- u$ cross-section derived previously

$$\frac{d\sigma^{\nu p}}{dy} = \frac{G_F^2}{\pi} \hat{s} d^P(x) dx$$

where \hat{s} is the centre-of-mass energy of the $\nu_\mu d$.

- Similarly for the \bar{u} contribution

$$\frac{d\sigma^{\nu p}}{dy} = \frac{G_F^2}{\pi} \hat{s} (1-y)^2 \bar{u}^p(x) dx.$$

- The neutrino-**proton** scattering cross section is obtained by summing the two contributions and using $\hat{s} = xs$ to give:

$$\frac{d^2\sigma^{\nu p}}{dx dy} = \frac{G_F^2}{\pi} sx \left[d^p(x) + (1-y)^2 \bar{u}^p(x) \right].$$

- The anti-neutrino **proton** differential cross section can be obtained in the same manner:

$$\frac{d^2\sigma^{\bar{\nu} p}}{dx dy} = \frac{GF^2}{\pi} sx \left[(1-y)^2 u^p(x) + \bar{d}^p(x) \right].$$

- For (anti)neutrino-**neutron** scattering:

$$\frac{d^2\sigma^{\nu n}}{dx dy} = \frac{G_F^2}{\pi} sx \left[d^n(x) + (1-y)^2 \bar{u}^n(x) \right]$$

$$\frac{d^2\sigma^{\bar{\nu} n}}{dx dy} = \frac{G_F^2}{\pi} sx \left[(1-y)^2 u^n(x) + \bar{d}^n(x) \right].$$

- As before, define neutron distributions functions in terms of those of the proton

$$u(x) \equiv u^p(x) = d^n(x); \quad d(x) \equiv d^p(x) = u^n(x)$$

$$\bar{u}(x) \equiv \bar{u}^p(x) = \bar{d}^n(x); \quad \bar{d}(x) \equiv \bar{d}^p(x) = \bar{u}^n(x)$$

$$\frac{d^2 \sigma^{\nu p}}{dx dy} = \frac{G_F^2}{\pi} s x \left[d(x) + (1-y)^2 \bar{u}(x) \right] \quad (129)$$

$$\frac{d^2 \sigma^{\bar{\nu} p}}{dx dy} = \frac{G_F^2}{\pi} s x \left[(1-y)^2 u(x) + \bar{d}(x) \right] \quad (130)$$

$$\frac{d^2 \sigma^{\nu n}}{dx dy} = \frac{G_F^2}{\pi} s x \left[u(x) + (1-y)^2 \bar{d}(x) \right] \quad (131)$$

$$\frac{d^2 \sigma^{\bar{\nu} n}}{dx dy} = \frac{G_F^2}{\pi} s x \left[(1-y)^2 d(x) + \bar{u}(x) \right] \quad (132)$$

- Because neutrino cross sections are very small, need massive detectors. These are usually made of iron, hence, one experimentally measure a combination of proton/neutron scattering cross sections.

- For an isoscalar target (i.e. equal numbers of protons and neutrons), the mean cross section per nucleon, N :

$$\frac{d^2\sigma^{\nu N}}{dx dy} = \frac{1}{2} \left(\frac{d^2\sigma^{\nu p}}{dx dy} + \frac{d^2\sigma^{\nu n}}{dx dy} \right)$$

$$\Rightarrow \boxed{\frac{d^2\sigma^{\nu N}}{dx dy} = \frac{G_F^2}{2\pi} s x \left[u(x) + d(x) + (1-y)^2(\bar{u}(x) + \bar{d}(x)) \right]}.$$

- Integrate over momentum fraction x :

$$\boxed{\frac{d\sigma^{\nu N}}{dy} = \frac{G_F^2}{2\pi} s \left[f_q + (1-y)^2 f_{\bar{q}} \right]} \quad (133)$$


where f_q and $f_{\bar{q}}$ are the total momentum fractions carried by the quarks and by the anti-quarks within a nucleon:

$$f_q \equiv f_d + f_u = \int_0^1 x [u(x) + d(x)] dx; \quad f_{\bar{q}} \equiv f_{\bar{d}} + f_{\bar{u}} = \int_0^1 x [\bar{u}(x) + \bar{d}(x)] dx.$$

- Similarly


$$\boxed{\frac{d\sigma^{\bar{\nu} N}}{dy} = \frac{G_F^2}{2\pi} s \left[(1-y)^2 f_q + f_{\bar{q}} \right]} \quad (134)$$

CDHS Experiment (CERN 1976-1984)




- 1250 tons
- Magnetized iron modules
- Separated by drift chambers


Study Neutrino Deep Inelastic Scattering



Experimental Signature:



CDHS (cont.)



- Measure energy of X , and call it E_X ,
- Measure muon momentum from curvature in B -field. and call it E_μ .
- For each event can then determine neutrino energy and y :

$$E_\nu = E_X + E_\mu$$

and then


$$E_\mu = (1 - y)E_\nu$$

implies







$$y = \left(1 - \frac{E_\mu}{E_\nu}\right).$$

Measured y Distribution

- CDHS measured y distribution
- Shapes can be understood in terms of (anti)neutrino – (anti)quark scattering



J. de Groot et al., Z.Phys. C1 (1979) 143



Measured Total Cross Sections


- Integrating the expressions for $\frac{d\sigma}{dy}$ (equations (133) and (134)):

$$\sigma^{\nu N} = \frac{G_F^2 s}{2\pi} \left[f_q + \frac{1}{3} f_{\bar{q}} \right]$$

and

$$\sigma^{\bar{\nu} N} = \frac{G_F^2 s}{2\pi} \left[\frac{1}{3} f_q + f_{\bar{q}} \right].$$


- In lab frame $p^\mu(\nu) = (E_\nu, 0, 0, +E_\nu)$ and $p^\mu(p) = (m_p, 0, 0, 0)$ so $s = (E_\nu + m_p)^2 - E_\nu^2 = 2E_\nu m_p + m_p^2 \approx 2E_\nu m_p$ therefore **the Neutrino DIS cross section is approximately proportional to lab. frame neutrino energy!**
- Measure cross sections can be used to determine fraction of protons momentum carried by quarks, f_q , and fraction carried by anti-quarks, $f_{\bar{q}}$.
- Find: $f_q \approx 0.41$; $f_{\bar{q}} \approx 0.08$.
- $\sim 50\%$ of momentum carried by gluons (which don't interact with virtual W -boson).
- If no anti-quarks in nucleons expect $\frac{\sigma^{\nu N}}{\sigma^{\bar{\nu} N}} = 3$.
- Including anti-quarks $\frac{\sigma^{\nu N}}{\sigma^{\bar{\nu} N}} \approx 2$.



Weak Neutral Current


- Neutrinos also interact via the Neutral Current. First observed in the Gargamelle bubble chamber in 1973. Interaction of muon neutrinos produce a final state muon.

$$\bar{\nu}_\mu + N \rightarrow \bar{\nu}_\mu + \text{hadrons}$$





F.J. Hasert et al., Phys. Lett. 46B (1973) 138

$$\bar{\nu}_\mu + e^- \rightarrow \bar{\nu}_\mu + e^-$$



F.J. Hasert et al., Phys. Lett. 46B (1973) 121

- Cannot be due to W exchange – this is first evidence for Z -boson!



Summary

- Derived neutrino/anti-neutrino – quark/anti-quark weak charged current (CC) interaction cross sections.
- Neutrino-nucleon scattering allows us to measure anti-quark content of protons and neutrons because:
 - ν couples to d and \bar{u} ,
 - $\bar{\nu}$ couples to u and \bar{d} ,
 - $\nu\bar{q}$ scattering is suppressed by factor $(1-y)^2$ compared with νq , and
 - $\bar{\nu}q$ scattering is suppressed by factor $(1-y)^2$ compared with $\bar{\nu}\bar{q}$.
- Further aspects of neutrino deep-inelastic scattering (expressed in general structure functions) are covered in Appendix XVII.
- Finally observe that neutrinos interact via weak neutral currents (NC).